

---

Subscribe

---

NEUROLOGICAL HEALTH

## How Dad's Stresses Get Passed Along to Offspring

Mouse studies show tiny intercellular pods convey to sperm a legacy of a father's hard knocks in life

By Esther Landhuis on November 8, 2018



Credit: Getty Images

---

ADVERTISEMENT

A stressed-out and traumatized father can leave scars in his children. New research suggests this happens because sperm “learn” paternal experiences via a mysterious mode of intercellular communication in which small blebs break off one cell and fuse with another.

Carrying proteins, lipids and nucleic acids, these particles ejected from a cell act like a postal system that extends to all parts of the body, releasing little packages known as extracellular vesicles. Their contents seem carefully chosen. “The cargo inside the vesicle determines not just where it came from but where it’s going and what it’s doing when it gets there,” says Tracy Bale, a neurobiologist at the University of Maryland School of Medicine.

Preliminary research Bale and others, announced this week at the annual meeting of the Society for Neuroscience in San Diego, shows how extracellular vesicles can regulate brain circuits and help diagnose neurodegenerative diseases—in addition to altering sperm to disrupt the brain health of resulting offspring.

---

ADVERTISEMENT

Striking evidence that harsh conditions affect a man's children came from crop failures and war ravaging Europe more than a century ago. In those unplanned human experiments, prolonged famine appeared to set off a host of health changes in future generations, including higher cholesterol levels and increased rates of obesity and diabetes. To probe the inheritance of such changes at the cellular level, Bale and co-workers performed a series of mouse experiments.

It is pretty easy to stress out a mouse. Stick one into a tube it cannot wriggle out of, soak its bedding or blast white noise—and stress hormone levels shoot up, much as they do in people worrying about finances or facing incessant pressure at work. Remarkably, the way a mouse physiologically responds to stress looks noticeably different if—months before conception—its father endured a period of stress. Somehow “their brain develops differently than if their dad hadn't experienced that stress,” says Chris Morgan, a postdoc in Bale's lab who helped create the mouse model.

The big question is how information about the paternal environment reaches the womb in the first place. After all, Morgan says, the “dad is only in there for one night, perhaps just a few hours.” Could his sperm carry memories of prior trauma? The idea seemed reasonable yet controversial. Because DNA is packed so tightly in the nucleus of a sperm cell, “the thought that [the cell] would respond to anything in the environment really boggled people's minds,” says Jennifer Chan, a former PhD student in Bale's lab who's now a postdoc at Icahn School of Medicine at Mount Sinai in New York City.

Rather, there must be some other kind of cell whose DNA does react to environmental changes—and that cell, she reasoned, could then relay that information to sperm cells to transmit at fertilization. She focused on a population of cells that interact with developing sperm by releasing molecules that help sperm grow and mature. They also secrete extracellular vesicles—and Chan showed it is these vesicles whose contents fuse with sperm cells, instilling memories of dad's prior stress.

In one set of experiments Chan stressed a group of male mice, let them mate and looked at stress responses in the pups. The clincher was a set of in vitro

fertilization–like experiments in which she collected sperm from a male mouse that had never experienced induced stress. Half his sperm went into a lab dish with vesicles previously exposed to stress hormones. The other half was cultured with vesicles that had no contact with stress hormones.



---

ADVERTISEMENT

Chan injected sperm cells from each batch into eggs from a nonstressed female, then implanted the fertilized eggs—zygotes—into the same foster mom. The pups from nonstressed zygotes developed normally. Pups from stress-exposed zygotes, however, showed the same abnormal stress response as those whose dads had experienced stress before mating. That showed extracellular vesicles act as the conduit for transmitting paternal stress signals to the offspring, Chan says.

The findings are “novel and of very high impact, especially when we consider the impact of military service or other work environments that can confer high stress,” says Robert Rissman, a neuroscientist at the University of California, San Diego, who was not involved with the research. “I think it would be important to better understand the specificity of the effect and how different types of stressors or strength of stressors can modulate this system.”

As a first step toward translating the findings to people, Morgan is collaborating with University of Pennsylvania psychiatrist Neill Epperson to track protein and RNA changes in human sperm samples. At the neuroscience meeting, Morgan presented data from a six-month study of 20 undergraduate and graduate students. Each month the participants came in and gave a sperm donation. They also completed a same-day survey asking how stressed they were feeling. Preliminary data suggests just several months after a student reports stress, his sperm shows changes in “small noncoding RNAs”—RNA molecules that do not get translated to protein but instead control which genes get turned on or off.



Sign up for *Scientific American's* newsletters.

Sign Up

---

Analyzing sperm from this group of healthy young men, the researchers plan to build a basic understanding of molecular changes linked with mild stresses such as taking final exams. In the future Bale and colleagues hope to compare these baseline fluctuations with changes induced by more prolonged life stressors such as post-traumatic stress disorder or neurological diseases such as autism and schizophrenia.

The molecular signatures in extracellular vesicles may also help researchers discover new ways to noninvasively diagnose or predict adverse health outcomes in offspring, says Gerlinde Metz, who studies transgenerational inheritance of stress responses at the University of Lethbridge in Alberta and was not involved with the research. If so, the vesicles could become the basis for a pioneering type of stress test.



---

ADVERTISEMENT

[Rights & Permissions](#)

---

## ABOUT THE AUTHOR(S)

### Esther Landhuis

Esther Landhuis is a freelance science journalist in the San Francisco Bay Area.

#### Recent Articles

Could Brain Stimulation Slow Cancer?

Experimental Huntington's Therapy Shows Promise in a Small Trial


Could the "Alzheimer's Gene" Finally Become a Drug Target?

**READ THIS NEXT****BEHAVIOR & SOCIETY****Separating Families May Cause Lifelong Health Damage**


June 20, 2018 — Sara Goudarzi

**BEHAVIOR & SOCIETY****Language Patterns Reveal Body's Hidden Response to Stress**


November 6, 2017 — Jo Marchant and Nature magazine

**BEHAVIOR & SOCIETY****Why Some People Are Born to Worry [Excerpt]**

May 5, 2017 — Daniel P. Keating

**MENTAL HEALTH****Traces of Genetic Trauma Can Be Tweaked**

April 15, 2017 — Erika Beras



## NEWSLETTER

Sign Up

---

## *From Genius to Madness*

Subscribe Now!

FOLLOW US

SCIENTIFIC AMERICAN ARABIC

العربية



[Store](#)

[About](#)

[Press Room](#)

[More](#)

Scientific American is part of Springer Nature, which owns or has commercial relations with thousands of scientific publications (many of them can be found at [www.springernature.com/us](http://www.springernature.com/us)). Scientific American maintains a strict policy of editorial independence in reporting developments in science to our readers.

© 2018 SCIENTIFIC AMERICAN, A DIVISION OF SPRINGER NATURE AMERICA, INC.

ALL RIGHTS RESERVED.